



## Pressure immobilization after North American Crotalinae snake envenomation

AMERICAN COLLEGE OF MEDICAL TOXICOLOGY, AMERICAN ACADEMY OF CLINICAL TOXICOLOGY, AMERICAN ASSOCIATION OF POISON CONTROL CENTERS, EUROPEAN ASSOCIATION OF POISON CONTROL CENTRES AND CLINICAL TOXICOLOGISTS, INTERNATIONAL SOCIETY ON TOXINOLOGY & ASIA PACIFIC ASSOCIATION OF MEDICAL TOXICOLOGY

**To cite this article:** AMERICAN COLLEGE OF MEDICAL TOXICOLOGY, AMERICAN ACADEMY OF CLINICAL TOXICOLOGY, AMERICAN ASSOCIATION OF POISON CONTROL CENTERS, EUROPEAN ASSOCIATION OF POISON CONTROL CENTRES AND CLINICAL TOXICOLOGISTS, INTERNATIONAL SOCIETY ON TOXINOLOGY & ASIA PACIFIC ASSOCIATION OF MEDICAL TOXICOLOGY (2011) Pressure immobilization after North American Crotalinae snake envenomation, *Clinical Toxicology*, 49:10, 881-882, DOI: [10.3109/15563650.2011.610802](https://doi.org/10.3109/15563650.2011.610802)

**To link to this article:** <http://dx.doi.org/10.3109/15563650.2011.610802>



Published online: 08 Nov 2011.



Submit your article to this journal [↗](#)



Article views: 552



View related articles [↗](#)



Citing articles: 1 View citing articles [↗](#)

POSITION STATEMENT

# Pressure immobilization after North American Crotalinae snake envenomation

AMERICAN COLLEGE OF MEDICAL TOXICOLOGY, AMERICAN ACADEMY OF CLINICAL TOXICOLOGY, AMERICAN ASSOCIATION OF POISON CONTROL CENTERS, EUROPEAN ASSOCIATION OF POISON CONTROL CENTRES AND CLINICAL TOXICOLOGISTS, INTERNATIONAL SOCIETY ON TOXINOLOGY, ASIA PACIFIC ASSOCIATION OF MEDICAL TOXICOLOGY

**Keywords** Snakes; pressure immobilization; Crotalinae; envenoming

## Background

The vast majority of venomous snake bites treated at health care facilities in the United States each year involve non-neurotoxic Crotalinae species.<sup>1</sup> Large case series reveal the major clinical effect associated with these envenomations to be local tissue injury. Extremity swelling and dermonecrosis are common, with compartment syndrome an infrequent but potentially limb-threatening effect of envenomation.<sup>2,3,4,5</sup> Life-threatening systemic toxicity and death are rare.

Historically, many first-aid measures have been employed in the treatment of snake bites, but none has been shown to improve patient outcome. Pressure immobilization is a technique routinely employed in the pre-hospital management of neurotoxic snake species in Australia. First described by Sutherland and colleagues in the 1970s, pressure immobilization involves wrapping the entire extremity with a bandage and then immobilizing the extremity with a splint.<sup>6</sup> The bandage should generate a pressure between 40–70 mm Hg in the upper extremity and 55–70 mmHg in the lower extremity in order to effectively delay systemic absorption of venom.<sup>7</sup>

Several animal studies have demonstrated delayed systemic absorption of venom with pressure immobilization.<sup>6,7,8</sup> However, studies have also revealed that pressure immobilization bandages are commonly applied incorrectly, even in a simulated setting following provider instructions and training.<sup>9,10,11,12</sup> Although the more common error is to apply the bandage too loosely, the bandage may function as a tourniquet when applied too tightly, causing limb ischemia, and may also increase systemic absorption of venom.<sup>7</sup>

Animal models of North American Crotalinae envenomation demonstrate delayed systemic absorption of venom and delayed mortality following application of pressure immobilization bandages.<sup>13,14,15</sup> However, the local effects of sequestering cytotoxic venom in the extremity are less clear. In a swine model of pressure immobilization following

*C atrox* lower extremity envenomation, intracompartmental pressure increased significantly compared to controls, from a non-surgical range to levels that would prompt fasciotomy.<sup>13</sup>

## Position

Given that the primary toxic effect of envenomation is local tissue injury, mortality is not an ideal outcome measure to extrapolate to human crotaline envenomation. Available evidence fails to establish the efficacy of pressure immobilization in humans, but indicates the possibility of serious adverse events arising from its use. The use of pressure immobilization for the pre-hospital treatment of North American Crotalinae envenomation is not recommended.

## Acknowledgements

The organizations acknowledge the efforts of Michael Levine, MD, and Michelle Ruha, MD in creating this position statement.

## Declaration of interest

The societies endorsing this position statement have no conflicts to report. This work has been endorsed by American College of Medical Toxicology, American Academy of Clinical Toxicology, American Association of Poison Control Centers, European Association of Poison Control Centres and Clinical Toxicologists, International Society of Toxicology and Asia Pacific Association of Medical Toxicology

## References

1. Bronstein AC, Spyker DA, Cantilena LR Jr, Green JL, Rumack BH, Giffin SL. 2009 Annual report of the American Association of Poison Control Centers' National Poison Data System (NPDS): 27th Annual Report. Clin Toxicol 2010; 48:979–1178.

2. Corneille MG, Larson S, Stewart RM, Dent D, Myers JG, Lopez PP, McFarland MJ, Cohn SM. A large single-center experience with treatment of patients with crotalid envenomations: outcomes with and evolution of antivenin therapy. *Am J Surg* 2006; 192(6):848–852.
3. Spiller HA, Bosse GM. Prospective study of morbidity associated with snakebite envenomation. *J Toxicol Clin Toxicol* 2003; 41(2): 125–130.
4. Tanen D, Ruha AM, Graeme K, Curry S. Epidemiology and hospital course of rattlesnake envenomations cared for at a tertiary referral center in central Arizona. *Acad Emerg Med* 2001; 8(2):177–182.
5. Thorson A, Lavonas EJ, Rouse AM, Kerns WP. Copperhead envenomations in the Carolinas. *J Toxicol Clin Toxicol* 2003; 41(1): 29–35.
6. Sutherland SK, Coulter AR, Harris RD. Rationalisation of first-aid measures for elapid snakebite. *Lancet* 1979; 27:183–185.
7. Howarth DM, Southee AE, Whyte IM. Lymphatic flow rates and first-aid in simulated peripheral snake or spider envenomation. *Med J Aust* 1994; 161:695–700.
8. German BT, Hack JB, Brewer K, Meggs WJ. Pressure-immobilization bandages delay toxicity in a porcine model of Eastern coral snake (*Micrurus fulvius fulvius*) envenomation. *Ann Emerg Med* 2005; 45: 603–608.
9. Canale E, Isbister GK, Currie BJ. Investigating pressure bandaging for snakebite in a simulated setting: bandage type, training, and the effect of transport. *Emerg Med Australas* 2009; 21:184–190.
10. Currie BJ, Canale E, Isbister GK. Effectiveness of pressure-immobilization first aid for snakebite requires further study. *Emerg Med Australas* 2008; 20:267–270.
11. Norris RL, Ngo J, Nolan K, Hooker G. Physicians and lay people are unable to apply pressure immobilization properly in a simulated snakebite scenario. *Wilderness Environ Med* 2005; 16:16–21.
12. Simpson ID, Tanwar PD, Andrade C, Kochar DK, Norris RL. The Ebbinghaus retention curve: training does not increase the ability to apply pressure immobilization in simulated snake bite – implications for snake bite first aid in the developing world. *Trans R Soc Trop Med Hyg* 2008; 102(5):451–459.
13. Bush SP, Green SM, Laack TA, Hayes WK, Cardwell MD, Tanen DA. Pressure immobilization delays mortality and increases intracompartmental pressure after artificial intramuscular rattlesnake envenomation in a porcine model. *Ann Emerg Med* 2004; 44:599–604.
14. Meggs WJ, Courtney C, O'Rourke D, Brewer KL. Pilot studies of pressure-immobilization bandages for rattlesnake envenomations. *Clin Toxicol* 2010; 48:61–63.
15. Sutherland SK, Coulter AR. Early management of bites by the Eastern diamondback rattlesnake (*Crotalus adamanteus*): studies in monkeys (*Macaca fascicularis*). *Am J Trop Med Hyg* 1981; 30: 497–200.